

Enhancement of the electric dipole moment of the electron in PbO

M. G. Kozlov*

Petersburg Nuclear Physics Institute, Gatchina, 188300, Russia

D. DeMille†

Physics Department, Yale University, New Haven, CT 06520

(Dated: February 2, 2008)

The $a(1)$ state of PbO can be used to measure the electric dipole moment of the electron d_e . We discuss a semiempirical model for this state, which yields an estimate of the effective electric field on the valence electrons in PbO. Our final result is an upper limit on the measurable energy shift, which is significantly larger than was anticipated earlier: $2|W_d|d_e \geq 2.4 \times 10^{25} \text{ Hz} \left[\frac{d_e}{e \text{ cm}} \right]$.

PACS numbers: 32.80.Ys, 11.30.Er, 31.30.Jv

In his pioneering work, Sandars pointed out that the effective electric field on a valence electron in a heavy atom is enhanced by a factor $\sim \alpha^2 Z^3$ relative to the applied laboratory field [1]. That started a long search for the electric dipole moment (EDM) of the electron d_e in atomic experiments ([2]). The most stringent limit on d_e follows from an experiment on atomic Tl ($Z=81$) [3].

Even larger enhancement is present in heavy polar diatomic molecules [2, 4, 5]. The heavy atom there is subjected to an internal E-field of $\sim 1 \text{ a.u.} \approx 5 \cdot 10^9 \text{ V/cm}$, which is further enhanced by the relativistic factor $\alpha^2 Z^3$. This effective field is many orders of magnitude larger than available laboratory fields; this makes diatomic molecules very attractive systems to look for d_e .

Since d_e is linked to the electron spin, one must work either with radicals, which have an unpaired electron in the ground state, or with excited states of “normal” molecules. Diatomic radicals with the ground state $\Sigma_{1/2}$ have large enhancement factors which can be relatively easily calculated [6, 7]. The first results of an EDM measurement in such a molecule (YbF) were recently published [8, 9]. The molecule PbO is a favorable candidate for a search for d_e in the excited state $a(1)$ [4, 10], and the group at Yale has begun EDM experiments on PbO [11]. It is therefore timely to estimate the effective internal field for the state $a(1)$ of PbO.

The interaction of d_e with an electric field $\mathbf{E}$ can be written in four-component Dirac notation as [12]:

$$H_d = 2d_e \begin{pmatrix} 0 & 0 \\ 0 & \boldsymbol{\sigma} \mathbf{E} \end{pmatrix}. \quad (1)$$

After averaging over the electronic wavefunction this interaction can be expressed in terms of an effective spin-rotational Hamiltonian [5, 6]:

$$H_d^{eff} = W_d d_e (\mathbf{J}_e \cdot \mathbf{n}), \quad (2)$$

where $\mathbf{J}_e$ is the electronic angular momentum and $\mathbf{n}$ is the unit vector along the molecular axis. In this paper we estimate W_d for the molecular state $a(1)$:

$$W_d \equiv d_e^{-1} \langle a(1) | H_d | a(1) \rangle, \quad (3)$$

where we used that $\langle a(1) | \mathbf{J}_e \cdot \mathbf{n} | a(1) \rangle \equiv \Omega[a(1)] = 1$. The Ω -doubling for states with $\Omega = 1$ is very small and even in a weak external electric field the energy eigenstates correspond to definite Ω rather than definite parity. The energy of the molecule can be then written as:

$$W(J, M, \Omega) = BJ(J+1) + W_d d_e \Omega - \frac{DE_0 \Omega M}{J(J+1)}, \quad (4)$$

where B is the rotational constant, E_0 is the external electric field, and D is the molecular dipole moment. The EDM contribution can be determined from the difference:

$$W(J, M, \Omega) - W(J, -M, -\Omega) = 2W_d d_e. \quad (5)$$

In order to estimate the matrix element in Eq. (3) we construct here a semiempirical wavefunction of the state $a(1)$. We use the MO LCAO approach, where each molecular orbital is expressed as a linear combination of atomic orbitals, and all molecular matrix elements are reduced to the sums of atomic matrix elements. The HFS or SO interactions as well as the EDM enhancement factor grow very rapidly with nuclear charge Z . Therefore, we are interested only in the Pb part of the MO LCAO expansion.

Analysis of the molecular observables requires knowledge of several atomic matrix elements for Pb. We calculate these in the Dirac-Fock approximation both for neutral Pb and for Pb^+ . Results are given in Table I for the orbitals $6s$ and $6p_j$. For the HFS operator we calculate the parameters $h_{k,k'}$ as defined in [5] (we use atomic units unless the opposite is stated explicitly):

$$h_{k,k'} = -\frac{g_n \alpha}{2m_p} \int_0^\infty (f_k g_{k'} + g_k f_{k'}) dr, \quad (6)$$

where $g_n = 0.59$ is the nuclear g-factor of ^{207}Pb , m_p is the proton mass, f_k and g_k are upper and lower components of the Dirac orbitals, and $k = (l-j)(2j+1)$ is the relativistic quantum number. For the EDM operator (1) in our minimal basis set there is only one nonzero radial

TABLE I: Atomic parameters $h_{k,k'}$ (GHz), w_{sp} (a.u.), and ξ (cm^{-1}), calculated in the Dirac-Fock approximation for Pb and Pb^+ . The relativistic quantum number k is equal to -1 , 1 , and -2 for $s_{1/2}$, $p_{1/2}$, and $p_{3/2}$ correspondingly.

	$h_{-1,-1}$	$h_{1,1}$	$h_{1,-2}$	$h_{-2,-2}$	w_{sp}	$\frac{w_{sp}}{\sqrt{-h_{-1,-1}h_{1,1}}}$	ξ
Pb^+	45.5	-8.9	-1.1	1.8	-34.1	-1.7	9452
Pb	42.3	-7.5	-0.9	1.4	-30.0	-1.7	8077
ratio	1.08	1.20	1.25	1.30	1.14	1.0	1.17

integral, between $6s_{1/2}$ and $6p_{1/2}$ orbitals:

$$w_{sp} = - \int_0^\infty g_{-1} g_1 \frac{d\phi}{dr} r^2 dr, \quad (7)$$

where ϕ is the atomic electrostatic potential. Finally, we also need the atomic SO constant ξ for the $6p$ shell:

$$H_{\text{SO}} = \xi \mathbf{l} \cdot \mathbf{s}, \quad \Rightarrow \quad \xi = \frac{2}{3} (\varepsilon_{6p_{3/2}} - \varepsilon_{6p_{1/2}}). \quad (8)$$

The last row of Table I gives the ratio of the radial integrals for the ion and for the atom. Note that the ratios are similar for all relevant integrals, and a simple relation between w_{sp} and HFS constants holds for both cases:

$$w_{sp} = -1.7 \sqrt{-h_{-1,-1} h_{1,1}}. \quad (9)$$

This relation is critical for semiempirical models of the EDM enhancement: it implies that the value of the EDM enhancement does not depend on what set of radial integrals (i.e. atomic or ionic) is used. Similarly (9) holds for other principal quantum numbers n , *e.g.* for $7s$ and $7p_j$. The particular choice of the radial integrals becomes important for the normalization of the wavefunction. The Pb atom in PbO is positively charged; therefore, below we use the ionic set of integrals from Table I.

In order to develop a semi-empirical model for the state $a(1)$, we have found it necessary to also consider the wavefunctions of several low-lying states of PbO . Previous work has shown that these states correspond to the configurations and nominal Λ, Σ -coupling terms as follows [13, 14]:

$$X(0^+)[^1\Sigma_0^+] : \sigma_1^2 \sigma_2^2 \sigma_3^2 \pi_1^4, \quad (10)$$

$$a(1)[^3\Sigma_1^+], C(0^+)[^3\Sigma_0^-], C'(1)[^3\Sigma_1^-] : \sigma_1^2 \sigma_2^2 \sigma_3^2 \pi_1^3 \pi_2, \quad (11)$$

$$A(0^+)[^3\Pi_0], B(1)[^3\Pi_1], D(1)[^1\Pi_1] : \sigma_1^2 \sigma_2^2 \sigma_3 \pi_1^4 \pi_2, \quad (12)$$

In the naive ionic model of PbO , the two $6p$ electrons from Pb move to O and close its $2p$ -shell. This suggests that the orbitals $\sigma_{1,2}$ and π_1 are centered on O, and that σ_3 is predominantly of the Pb $6s$ -type. However, below we do not impose any constraints on the MO LCAO coefficients for these molecular orbitals, based on this expectation. Note that only the orbitals σ_3 and $\pi_{1,2}$ contribute

to the spin-density of the molecular states under consideration. Thus, we do not need to know the orbitals $\sigma_{1,2}$, and below we omit the index for the orbital σ_3 .

Now we specify coefficients of the MO LCAO expansion for the three valence orbitals of interest:

$$|\sigma_\omega\rangle = S_s |6s_{1/2,\omega}\rangle + S_p \left(-2\omega \sqrt{\frac{1}{3}} |6p_{1/2,\omega}\rangle + \sqrt{\frac{2}{3}} |6p_{3/2,\omega}\rangle \right), \quad (13a)$$

$$|\pi_{i,\omega}\rangle = P_i \left(2\omega \sqrt{\frac{2}{3}} |6p_{1/2,\omega}\rangle + \sqrt{\frac{1}{3}} |6p_{3/2,\omega}\rangle \right), \quad (13b)$$

$$|\pi_{i,\omega'}\rangle = P_i |6p_{3/2,\omega'}\rangle, \quad (13c)$$

where $\omega = \pm 1/2$ and $\omega' = \pm 3/2$. The numerical coefficients are chosen to account for the quantum number λ : $\lambda = 0(1)$ for $\sigma(\pi)$ orbitals. In order to calculate W_d we must determine the 4 parameters in Eqs. (13). Below we try to constrain these parameters using experimental information about states (11) and (12). To simplify the notation, we define the ground state of the molecule as vacuum. Then each of the excited states in Eqs. (11) and (12) is a two-particle state with one hole and one electron. We do not use any special notation for the hole states; instead we simply write the hole orbital in front of the electron one. We construct wavefunctions of these states from the orbitals (13), using at the first stage the Λ, Σ -coupling scheme classification:

$$\begin{aligned} |a(1)[^3\Sigma_1^+]\rangle &= \frac{1}{\sqrt{2}} (|\pi_{1,\lambda=-1} \pi_{2,\lambda=1}\rangle + |\pi_{1,\lambda=1} \pi_{2,\lambda=-1}\rangle) |\uparrow\uparrow\rangle \\ &= \frac{1}{\sqrt{2}} (|\pi_{1,-1/2} \pi_{2,3/2}\rangle + |\pi_{1,3/2} \pi_{2,-1/2}\rangle); \end{aligned} \quad (14a)$$

$$\begin{aligned} |A(0^+)[^3\Pi_0]\rangle &= \frac{1}{\sqrt{2}} (|\sigma \pi_{2,\lambda=1}\rangle |\downarrow\downarrow\rangle - |\sigma \pi_{2,\lambda=-1}\rangle |\uparrow\uparrow\rangle) \\ &= \frac{1}{\sqrt{2}} (|\sigma_{-1/2} \pi_{2,1/2}\rangle - |\sigma_{1/2} \pi_{2,-1/2}\rangle); \text{ etc.} \end{aligned} \quad (14b)$$

We write each wavefunction in both λ - σ and ω - ω representations; the latter is more convenient for our purposes.

The rules for calculating hole matrix elements follow from the fact that the hole in the state $|\omega\rangle$ actually means the absence of the electron in the state $|- \omega\rangle$. Thus, the expectation value for an electronic operator $\hat{P}$ over the hole state $|\omega\rangle$ can be written as:

$$\langle \omega | \hat{P} | \omega \rangle_h \equiv - \langle -\omega | \hat{P} | -\omega \rangle_e \stackrel{T}{=} \mp \langle \omega | \hat{P} | \omega \rangle_e, \quad (15)$$

where we applied the time-reversal operation T . Thus the final sign depends on the time-reversal symmetry of $\hat{P}$, with the minus sign corresponding to a T -even electronic operator. For example, the HFS interaction is given by the product of the T -odd electronic vector $\hat{\mathbf{A}} \mathbf{J}_e$ and the nuclear spin $\mathbf{I}$. Thus, for the HFS interaction the plus

sign in Eq. (15) is correct. A similar argument shows that the SO constant ξ for a hole has the opposite sign as for an electron.

From Eqs. (14), the first-order SO splitting Δ_{AB} between states $A(0^+)$ and $B(1)$ is:

$$\Delta_{AB} = \frac{\xi}{2} [\langle \pi_{2,3/2} | \mathbf{l} s | \pi_{2,3/2} \rangle - \langle \pi_{2,1/2} | \mathbf{l} s | \pi_{2,1/2} \rangle] = \frac{\xi P_2^2}{2}. \quad (16)$$

Using the experimental value of this splitting [15] and the ionic value for ξ from Table I, we estimate P_2 :

$$P_2^2 = \frac{2\Delta_{AB}}{\xi} = \frac{2 \cdot 2420}{9450} = 0.51. \quad (17)$$

We see that the orbital π_2 has a large contribution from the Pb orbital $6p$. The data on energy levels [15] shows then that for all levels with one electron in the π_2 orbital, the SO interaction is comparable to the splittings between these levels. Therefore, there must be significant SO mixing between such states.

We start with the mixing within configuration $\sigma^2\pi_1^3\pi_2$. The mixing angle α between states $a(1)$ and $C'(1)$ is:

$$\alpha \approx \frac{\langle {}^3\Sigma_1^- | H_{\text{so}} | {}^3\Sigma_1^+ \rangle}{|\Delta_{aC'}|} = \frac{\xi_1 + \xi_2}{2|\Delta_{aC'}|}, \quad (18)$$

where $\Delta_{aC'}$ is the energy splitting between $a(1)$ and $C'(1)$, and $\xi_i \equiv \xi P_i^2$. If we assume that $P_1^2 \ll P_2^2$ (corresponding to the naive ionic model), we can estimate the value of α :

$$\alpha \approx \frac{\xi_2}{2|\Delta_{aC'}|} \approx 0.3, \quad (19)$$

and write the new wavefunction in the form:

$$|a(1)\rangle = c_\alpha |\pi_{1,3/2}\pi_{2,-1/2}\rangle + s_\alpha |\pi_{1,-1/2}\pi_{2,3/2}\rangle, \quad (20)$$

$$c_\alpha \equiv \cos\left(\frac{\pi}{4} - \alpha\right), \quad s_\alpha \equiv \sin\left(\frac{\pi}{4} - \alpha\right). \quad (21)$$

SO interaction also mixes configuration $\sigma^2\pi_1^3\pi_2$ with configurations $\sigma\pi_1^3\pi_2^2$ and $\sigma\pi_1^4\pi_2$. These mixings can be accounted for by substitution of the original orbitals $|\pi_{i,1/2}\rangle$ with the perturbed orbitals

$$|\tilde{\pi}_{i,1/2}\rangle = c_i |\pi_{i,1/2}\rangle + s_i |\sigma_{1/2}\rangle. \quad (22)$$

There is no experimental information about levels of the configuration $\sigma\pi_1^3\pi_2^2$, so we cannot reliably estimate the mixing parameter s_2 . In contrast, both levels with $\Omega = 1$ of the configuration $\sigma\pi_1^4\pi_2$ are known [i.e., B(1) and D(1)]. That allows us to write for s_1 the estimate:

$$s_1 = 2.8 s_\alpha^2 P_1 S_p. \quad (23)$$

These SO mixings then lead to the final form of the wavefunction of the state $a(1)$:

$$|a(1)\rangle = c_\alpha |\pi_{1,3/2}\tilde{\pi}_{2,-1/2}\rangle + s_\alpha |\tilde{\pi}_{1,-1/2}\pi_{2,3/2}\rangle. \quad (24)$$

The G -factor for the state (24) is given by:

$$G_{\parallel} = \langle a(1) | L_0 + 2S_0 | a(1) \rangle = 2 - s_\alpha^2 s_1^2 - c_\alpha^2 s_2^2. \quad (25)$$

The measured value $G_{\parallel} = 1.84(3)$ [16] corresponds to the following equation for mixing parameters:

$$s_\alpha^2 s_1^2 + c_\alpha^2 s_2^2 = 0.16(3). \quad (26)$$

The signs of the parameters $s_{1,2}$ should be chosen so that the contribution of atomic orbital $6p_{1/2}$ to the molecular orbital σ is increased: in this case relativistic corrections to the binding energy of the σ orbital are positive.

The matrix element of the HFS interaction for the state $a(1)$ (24) has the form:

$$\begin{aligned} \langle a(1) | H_{\text{hfs}} | a(1) \rangle &= \\ &= c_\alpha^2 [\langle \pi_{1,3/2} | h_{\text{hfs}} | \pi_{1,3/2} \rangle - \langle \tilde{\pi}_{2,1/2} | h_{\text{hfs}} | \tilde{\pi}_{2,1/2} \rangle] \\ &+ s_\alpha^2 [\langle \pi_{2,3/2} | h_{\text{hfs}} | \pi_{2,3/2} \rangle - \langle \tilde{\pi}_{1,1/2} | h_{\text{hfs}} | \tilde{\pi}_{1,1/2} \rangle]. \end{aligned} \quad (27)$$

We use expressions from Ref. [5] for the one-electron matrix elements and numbers from Table I, combined with the measurement of the hyperfine constant for the state $a(1)$: $A_{\parallel} = -4.1$ GHz [16], to find another equation relating the various coefficients of the model:

$$\begin{aligned} &30(c_\alpha^2 s_2^2 + s_\alpha^2 s_1^2) S_s^2 + 1.8(c_\alpha^2 s_2^2 + s_\alpha^2 s_1^2) S_p^2 \\ &+ (4.6s_\alpha^2 c_1^2 - 1.4c_\alpha^2) P_1^2 + (4.6c_\alpha^2 c_2^2 - 1.4s_\alpha^2) P_2^2 \\ &- 4.7s_\alpha^2 c_1 s_1 P_1 S_p - 4.7c_\alpha^2 c_2 s_2 P_2 S_p = 4.1. \end{aligned} \quad (28)$$

(Note that the formulae of Ref. [5] are strictly applicable only for orbitals and states with $\omega = \Omega = 1/2$. Eq. (28) takes into account simple modifications of these formulae for the present situation.)

We now have five equations, namely: (17), (19), (23), (26), and (28) on seven parameters of the wavefunction (24). That leaves us with two independent parameters of the model. We introduce two additional constraints, which account for normalization and the Pauli principle:

$$S_s^2 + S_p^2 \leq N_0, \quad P_1^2 + P_2^2 \leq N_0. \quad (29)$$

We choose $N_0 = 1.2$ here in order to account for inaccuracy of the Hartree-Fock approximation used to determine the atomic parameters in Table I.

The parameters α and P_2 are unambiguously fixed by Eqs. (17) and (19). We choose s_1 and P_1 as free parameters and solve Eqs. (23), (26), and (28) for parameters s_2 , S_p and S_s . After that we reject solutions which do not meet the constraints (29). A typical solution is:

$$\begin{cases} \alpha = 0.3; & P_2 = 0.714; \\ s_1 = 0.107; & P_1 = 0.503; \\ s_2 = 0.449; & S_p = 0.349; & S_s = 0.812. \end{cases} \quad (30)$$

Some of the parameters appear relatively well-defined, while others are not. The variation ranges are:

$$\begin{cases} s_1 \leq 0.2; & 0.4 \leq s_2 \leq 0.5 \\ S_p^2 \leq 0.5; & S_s^2 \geq 0.5. \end{cases} \quad (31)$$

TABLE II: Dependence of the EDM constant W_d (in a.u.) on the parameters of the model.

$A_{ }$ (GHz)	$G_{ }$	α	P_2^2	$ W_d $	
				max	min
-4.1	1.84	0.30	0.51	19.6	13.7
-4.1	1.81	0.30	0.51	19.1	12.0
-4.1	1.87	0.30	0.51	20.3	15.4
-4.1	1.84	0.24	0.51	19.8	13.2
-4.1	1.84	0.36	0.51	20.3	15.5
-4.1	1.84	0.30	0.41	18.2	12.1
-4.1	1.84	0.30	0.61	20.3	15.1
-3.3	1.84	0.30	0.51	17.0	11.2
-4.9	1.84	0.30	0.51	22.0	16.4

The parameter P_1 appears to be restricted only by the normalization condition (29).

It may be possible to add some restrictions to reduce the ranges of variation in Eq. (31). For example, the relatively large value of s_2 should require a large value of S_p . However, such additional restrictions would add arbitrariness to the model and may affect its reliability. We use only the minimal set of constraints to determine the range of possible values of W_d .

For the wavefunction (24), there are two contributions to the EDM parameter W_d from each of the one-electron orbitals with $|\omega| = 1/2$:

$$W_d = -c_\alpha^2 W_d^{\tilde{\pi}_2} - s_\alpha^2 W_d^{\tilde{\pi}_1} \quad (32a)$$

$$= \frac{4w_{sp}}{\sqrt{3}} S_s \left(\sqrt{2} c_\alpha^2 c_2 s_2 P_2 - c_\alpha^2 s_2^2 S_p \right. \\ \left. + \sqrt{2} s_\alpha^2 c_1 s_1 P_1 - s_\alpha^2 s_1^2 S_p \right). \quad (32b)$$

We find that the first term in (32b) always dominates the sum. The second term is not negligible, but the final two terms contribute $\lesssim 10\%$. It is important that the leading contribution to W_d is similar to the first term in Eq. (28), which dominates the HFS. This implies that the parameter W_d is well-constrained even though some of the parameters of the wavefunction are not. We obtain:

$$|W_d| = 16.6 \pm 3.0 \text{ a.u.}, \quad (33)$$

where the uncertainty reflects the range of values found within the model just described.

It is also important to check how W_d depends on the “fixed” parameters α and P_2 , as well as on the input data for $A_{||}$ and $G_{||}$, since our model relating the MO LCAO coefficients to these parameters is rather crude. In Table II we solve the model equations for values of these quantities varying from the best values by $\pm 20\%$. We find that this variation of the input parameters widens the range for W_d substantially (to ± 5.4 a.u.), but still does not allow dramatically smaller values of W_d .

It is known from previous calculations of W_d for other diatomic molecules, that correlation corrections tend to

decrease the result by 10–20% from the Hartree-Fock level. Therefore, we state our final result as a conservative lower limit on W_d :

$$|W_d| \geq 10 \text{ a.u.} = 12 \cdot 10^{24} \frac{\text{Hz}}{e \text{ cm}}. \quad (34)$$

This lower bound is several times larger than earlier, naive estimates which did not consider the effect of SO mixing on the (nominally) π -type orbitals of the $a(1)$ state [5]. Our model shows significant similarity between the orbital $\tilde{\pi}_{2,1/2}$ in PbO and the single valence orbital in the ground state of the free radical PbF. It is thus natural that our bound is close to the value calculated for PbF [21]. (Coincidentally, our bound is also similar to the calculated value for YbF [17, 18, 19, 20].) However, we stress that this first semiempirical estimate of the effective field in PbO has very limited accuracy. Thus, more elaborate calculations of the $a(1)$ state are highly desirable.

MK thanks the University of New South Wales for hospitality and acknowledges support from RFBR, grant No 02-02-16387. DD was supported by NSF Grant PHY9987846, a NIST Precision Measurement Grant, Research Corporation, the David and Lucile Packard Foundation, and the Alfred P. Sloan Foundation.

* Electronic address: mgk@MF1309.spb.edu

† Electronic address: david.demille@yale.edu

- [1] P. G. H. Sandars, Phys. Lett. **14**, 194 (1965).
- [2] E. D. Commins, Adv. Atom. Mol. Opt. Phys. **40**, 1 (1999).
- [3] B. C. Regan *et al.*, Phys. Rev. Lett. **88**, 071805 (2002).
- [4] O. P. Sushkov and V. V. Flambaum, Sov. Phys.-JETP **48**, 608 (1978).
- [5] M. Kozlov and L. Labzowski, J. Phys. B **28**, 1931 (1995).
- [6] M. G. Kozlov, Sov. Phys.-JETP **62**, 1114 (1985).
- [7] M. G. Kozlov and V. F. Ezhov, Phys. Rev. A **49**, 4502 (1994).
- [8] E. A. Hinds, Phys. Scr. **T70**, 34 (1997).
- [9] J. J. Hudson *et al.*, e-print:hep-ex/0202014.
- [10] D. DeMille *et al.*, Phys. Rev. A **61**, 052507 (2000).
- [11] D. DeMille *et al.*, AIP Conference Proceedings (AIP, Melville, New-York, 2001), vol. 596, pp. 72–83.
- [12] I. B. Khriplovich, *Parity non-conservation in atomic phenomena* (Gordon and Breach, New York, 1991).
- [13] R. C. Oldenberg, C. R. Dickson, and R. N. Zare, J. Mol. Spectrosc. **58**, 283 (1975).
- [14] K. Balasubramanian and K. S. Pitzer, J. Phys. Chem. **87**, 4857 (1983).
- [15] K. P. Huber and G. Herzberg, *Constants of Diatomic Molecules* (Van Nostrand, New York, 1979).
- [16] L. R. Hunter *et al.*, Phys. Rev. A **65**, 030501 (2002).
- [17] M. G. Kozlov, J. Phys. B **30**, L607 (1997).
- [18] F. Parpia, J. Phys. B **31**, 1409 (1998).
- [19] H. M. Quiney, H. Skaane, and I. P. Grant, J. Phys. B **31**, L85 (1998).
- [20] N. S. Mosyagin, M. G. Kozlov, and A. V. Titov, J. Phys. B **31**, L763 (1998).

- [21] M. G. Kozlov *et al.*, J. Phys. B **20**, 4939 (1987).